

## DETERMINATION OF 17 $\alpha$ -HYDROXYLASE-C<sub>17,20</sub>-LYASE (P450<sub>17 $\alpha$</sub> ) ENZYME ACTIVITIES AND THEIR INHIBITION BY SELECTED STEROIDAL PICOLYL AND PICOLINYLIDENE COMPOUNDS

NIKOLETTA SZABÓ,<sup>1</sup> JOVANA J. AJDUKOVIĆ,<sup>2</sup> EVGENIJA A. DJURENDIĆ,<sup>2</sup>  
MARIJA N. SAKAČ,<sup>2</sup> IMRE IGNÁTH,<sup>1</sup> JÁNOS GARDI,<sup>1</sup> GÁBOR MAHMOUD,<sup>1</sup>  
OLIVERA R. KLISURIĆ,<sup>3</sup> SUZANA JOVANOVIĆ-ŠANTA,<sup>2</sup>  
KATARINA M. PENOV GAŠI<sup>2</sup> and MIHÁLY SZÉCSI<sup>1</sup> \*

<sup>1</sup>1<sup>st</sup> Department of Medicine, University of Szeged, Korányi fasor 8–10, H-6720 Szeged, Hungary

<sup>2</sup>Department of Chemistry, Biochemistry and Environmental Protection, Faculty of Sciences, University of Novi Sad, 3 Trg D. Obradovića, 21000 Novi Sad, Serbia

<sup>3</sup>Department of Physics, Faculty of Sciences, University of Novi Sad, 4 Trg D. Obradovića, 21000 Novi Sad, Serbia

(Received: April 2, 2014; accepted: May 14, 2014)

17 $\alpha$ -hydroxylase-C<sub>17,20</sub>-lyase (P450<sub>17 $\alpha$</sub> ) is a key regulator enzyme of the steroid hormone biosynthesis in both the adrenals and the testes. Inhibition of this enzyme can block androgen synthesis in an early step, and may thereby be useful in the treatment of several androgen-dependent diseases. We developed radio-substrate *in vitro* incubation methods for the determination of the distinct 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase activities of the enzyme using rat testicular homogenate as enzyme source. With this method we have studied the inhibiting activity of selected steroidal picolyl and picolinylidene compounds. Tests revealed a substantial inhibitory action of the 17-picolinyliden-androst-4-en-3-one compound.

**Keywords:** EC 1.14.99.9 / EC 4.1.2.30 – 17 $\alpha$ -hydroxylase-C<sub>17,20</sub>-lyase (P450<sub>17 $\alpha$</sub> ) inhibitors – *exo*-heterocyclic steroids – rat testis – abiraterone

### INTRODUCTION

17 $\alpha$ -hydroxylase-C<sub>17,20</sub>-lyase (P450<sub>17 $\alpha$</sub> ) plays an important role in the pathways of steroid hormone biosynthesis [26, 28]. This is a cytochrome P450 enzyme located on the endoplasmic reticulum of the cells. It is expressed and active in the adrenal glands, testes, ovaries and placenta, primarily. The P450<sub>17 $\alpha$</sub>  is capable of catalysing two reactions engaged on a single active site and is active both in the  $\Delta^5$  and  $\Delta^4$  pathways. The enzyme converts C<sub>21</sub> precursors pregnenolone and progesterone (Prog) first to their 17 $\alpha$ -hydroxy derivatives 17-hydroxypregnenolone and 17-hydroxyprogesterone (17OHProg). A subsequent side chain cleavage process forms C<sub>19</sub> steroids, dehydroepiandrosterone (DHEA) and androstenedione (4-en-dione) (Fig. 1). Prog is a precursor of mineralocorticoids, whereas its hydroxylated derivative 17OHProg may result in glucocorticoids. C<sub>19</sub> steroids are androgen prehormones: in steroido-

\*Corresponding author; e-mail: szecsi.mihaly@med.u-szeged.hu

genic tissues and target organs they can be further metabolized to more potent androgens such as testosterone and dihydrotestosterone. Inhibition of P450<sub>17 $\alpha$</sub>  can block androgen synthesis, and may be competent in pharmacological treatment of hormone dependent disorders [4, 16, 18]. This enzyme is also a potential target for a drug designed to treat androgen dependent prostatic carcinoma [12, 27]. Inhibitory action was found to be related to lone electron pair coordination to the heme iron atom at the active site of the enzyme and many steroidal compounds bearing different heteroaromatic moieties on ring D of the sterane skeleton were analyzed against P450<sub>17 $\alpha$</sub> . Pyridyl analogues, for example, exhibited highly effective inhibition of the enzyme, and abirateron [17-(3-pyridyl)androsta-5,16-dien-3 $\beta$ -ol] is recently introduced into medical application [2, 32, 36].

We report here radiosubstrate *in vitro* incubation methods for the specific determination of 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase activities. Using these methods we have studied inhibitory action of selected steroidal 17-picolyl and 17-picolinylidene derivatives exerted towards the rat P450<sub>17 $\alpha$</sub> . Earlier publications describe synthesis methods and chemical properties of these compounds [7, 14, 25].




Fig. 1. Steroid biotransformations by 17 $\alpha$ -hydroxylase-C<sub>17,20</sub>-lyase (P450<sub>17 $\alpha$</sub> )

## MATERIALS AND METHODS

### *Steroid standards and other chemicals*

Radioactive [4-<sup>14</sup>C]Prog with specific activity (S.A.) of 46 mCi/mmol and [1,2,6,7-<sup>3</sup>H]17OH-Prog, S. A. = 50 Ci/mmol was purchased from the Radiochemical Centre, Amersham International (Amersham, UK) and American Radiolabeled Chemicals (St. Louis, MO, USA), respectively. Non-radioactive Prog and 17OH-Prog and 4-en-dione standards and reference inhibitor ketoconazole were obtained from Sigma (St. Louis, MO, USA) and Fluka (Buchs, Switzerland). Abiraterone was prepared with the synthesis method of Potter et al. [33] and was provided by Professor Gyula Schneider and Professor János Wölfling (Department of Organic Chemistry, University of Szeged). Other chemicals and solvents with purity of analytical grade were purchased from Sigma (St. Louis, MO, USA).

### *Preparation of rat testicular homogenates*

Testicular tissue was obtained from male rats (12-week old, 200–250 g body mass) via surgeries under ether anaesthesia. Tissue specimens were washed with an isotonic solution of NaCl. 1.0 g pieces of the testicles of five animals were mixed and the tissue sample was homogenized with an Ultra-Turrax in 20.0 ml 0.1 M HEPES buffer (pH = 7.3) containing 1 mM EDTA and 1 mM dithiothreitol. The homogenate was centrifuged at 1500 g for 10 min at 4 °C. The supernatant was divided into portions for storage at –70 °C.

### *In vitro incubations*

17 $\alpha$ -hydroxylase: Testis homogenate aliquots containing suitable amount of enzyme (equivalent to 10 mg of testicular tissue) were incubated with 1  $\mu$ M [<sup>14</sup>C]Prog (25,000 dpm) in the presence of 1 mM NADPH in total volume of 200  $\mu$ l HEPES buffer incubation medium (pH = 7.3). [<sup>14</sup>C]Prog was added to the incubate in 20  $\mu$ l 30 v/v% propylene glycol/HEPES buffer solution. Incubation was carried out at 37 °C for 20 min in air. The enzymatic reaction was stopped by the addition of ethyl acetate and freezing. After extraction, unlabeled carriers of Prog, 17OH-Prog and 4-en-dione were added to the samples to aid visualization of steroid spots on the chromatography plate by ultraviolet light. The three steroids were separated by thin layer chromatography (TLC) on Kieselgel-G (Merck Si 254 F) layers (0.25 mm thick) with the solvent system dichloromethane/diisopropyl ether/ethyl acetate (75:15:10 v/v). Ultraviolet light was used to trace the separated steroids. Prog, 17OH-Prog and 4-en-dione were found at retention factors 0.67, 0.24 and 0.48, respectively. Spots were cut out and the radioactivity of the 17OH-Prog and 4-en-

dione formed, and the Prog remaining was measured by liquid scintillation counting (Packard Tri-Carb 2200CA). The  $17\alpha$ -hydroxylase activity was calculated from the conversions corrected with recoveries of the  $17\text{OH-Prog}$  and 4-en-dione formed and expressed in picomoles.

$C_{17,20}$ -lyase: Testis homogenate aliquots containing suitable amount of the enzyme (equivalent to 30 mg of testicular tissue) were incubated with  $1\ \mu\text{M}$  [ $^3\text{H}$ ]17OH-Prog (300,000 dpm) in the presence of 1 mM NADPH in HEPES buffer incubation medium for 20 min at  $37\ ^\circ\text{C}$ . The substrate, in this procedure, was prepared with mix of the tritiated tracer and the non-labelled 17OH-Prog, and was added to the incubate also in  $20\ \mu\text{l}$  30 v/v% propylene glycol/HEPES buffer solution. Work-up procedures of the incubates and isolation of the 4-en-dione product and the substrate 17OH-Prog rest were performed as described above. The  $C_{17,20}$ -lyase activity was calculated from the conversion corrected with recovery of 4-ene-dione formed.




Fig. 2. Concentration dependent inhibition of  $17\alpha$ -hydroxylase and  $C_{17,20}$ -lyase activities of the  $\text{P450}_{17\alpha}$  and logit-log transformation with linear regression analysis of inhibition data for the determination of  $\text{IC}_{50}$  values. P: enzymatic conversion in the presence of inhibitor,  $P_0$ : enzymatic conversion of the control incubation in the absence of inhibitor. Relative conversion =  $P/P_0$ ,  $\text{Logit}(P) = \ln[P/(P_0 - P)]$ .  $\text{IC}_{50}$  values are derived from intercepts on the axis of  $\ln$  (inhibitor concentration), as 50% inhibition results in logit parameter equal to zero

### *Inhibitor tests*

The inhibitory effect of the new test compounds was investigated at 0.01–50  $\mu\text{M}$  concentration interval. They were added to the incubate in dimethylsulfoxide solution in a maximal volume of 10  $\mu\text{l}$ . Measurements were performed at least in four to six different concentrations and at least two experiments were done with each concentration. The control experiments were performed without the test substances in every incubation series. Abiraterone and a non-steroidal compound ketoconazole, known potent P450<sub>17 $\alpha$</sub>  inhibitors were used as reference compounds. The inhibitory effects of the compounds investigated in our present work are given in terms of IC<sub>50</sub> values, i.e. the concentration of inhibitor at which the 17 $\alpha$ -hydroxylase or C<sub>17,20</sub>-lyase activity was decreased to 50%. IC<sub>50</sub> results were calculated by linear regression analysis following a logit-log transformation of the data (Fig. 2), and the standard deviations were determined from the fitted lines. Compounds exerting weaker inhibitory effect were investigated at 50  $\mu\text{M}$  concentration and relative conversions compared to control incubates were determined in these cases. Two experiments were performed and mean values and standard deviations were calculated.

## RESULTS

### *Methodological investigations*

The optimum conditions for the measurement of 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase activities in rat testicular homogenate were determined in preliminary experiments. Enzyme activities were then measured under conditions of linearity with respect to enzyme concentration, and incubation time. Conversion of the substrate in non-inhibited control incubates was set to 10–15%. We achieved similar turnovers and optimal conversion rates applying higher enzyme concentration for the rate limiting C<sub>17,20</sub>-lyase reaction than for 17 $\alpha$ -hydroxylase activity, under the same incubation conditions. In 17 $\alpha$ -hydroxylase method we have found that 15–25% of the 17OH-Prog product was consecutively converted to 4-en-dione. This rate was independent from incubation conditions (time, substrate or enzyme concentrations) as the cleavage process takes place without dissociation of 17OH-Prog from the active centre of P450<sub>17 $\alpha$</sub> . Dimethylsulfoxide and propylene glycol were present in the incubates with 5.0 and 3.0 v/v% concentrations, respectively. These organic solvent contents did not reduce the enzyme activities substantially. Procedural loss of the isolation steps was determined by regained radioactivity and the recovery was found around 70% for Prog, 60% for 17OH-Prog, and 80% for 4-en-dione. Coefficients of variation of enzyme activity results in repeated measurements were within  $\pm 10\%$ . IC<sub>50</sub> parameters of references ketoconazole and abiraterone were found  $1.86 \pm 0.25 \mu\text{M}$  and  $0.034 \pm 0.003 \mu\text{M}$  for 17 $\alpha$ -hydroxylase, and  $0.32 \pm 0.02 \mu\text{M}$  and  $0.0125 \pm 0.0015 \mu\text{M}$  for C<sub>17,20</sub>-lyase in our P450<sub>17 $\alpha$</sub>  inhibition test, respectively. These reference results are in good agreement with values found in the literature [4, 9, 15, 18, 23, 31–32, 34].

*Table 1*  
*In vitro* inhibition of 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase activities of the rat testicular P450<sub>17 $\alpha$</sub>  enzyme by picolyl (1–3) and picolinylidene (4–6) androstene compounds







Structural formula, name and number of compound	17 $\alpha$ -Hydroxylase inhibition		C <sub>17,20</sub> -Lyase inhibition	
	Relative conversion at 50 $\mu$ M [%]	IC <sub>50</sub> [ $\mu$ M]	Relative conversion at 50 $\mu$ M [%]	IC <sub>50</sub> [ $\mu$ M]
 17 $\alpha$ -picolyl-androst-5-en-3 $\beta$ ,17 $\beta$ -diol (1)	95 $\pm$ 0.6	–	NI	–
 17 $\alpha$ -picolyl-androst-4-en-3 $\beta$ ,17 $\beta$ -diol (2)	90 $\pm$ 4	–	66 $\pm$ 13	–
 17 $\beta$ -hydroxy-17 $\alpha$ -picolyl-androst-4-en-3-on (3)	87 $\pm$ 10	–	73 $\pm$ 11	–

Table 1 (cont.)

Structural formula, name and number of compound	17α-Hydroxylase inhibition		C <sub>17,20</sub> -Lyase inhibition	
	Relative conversion at 50 μM [%]	IC <sub>50</sub> [μM]	Relative conversion at 50 μM [%]	IC <sub>50</sub> [μM]
 17( <i>E</i> )-picolinyliden-androst-5-en-3β-ol ( <b>4</b> )	45±4	38±1.4	–	8.2±1.8
 17( <i>E</i> )-picolinyliden-androst-4-en-3β-ol ( <b>5</b> )	48±2	48±1.7	23±1	14±1.2
 17( <i>E</i> )-picolinyliden-androst-4-en-3-on ( <b>6</b> )	–	5.9±1.2	–	2.5±1.4

Results of IC<sub>50</sub> ± S.D. in μM and relative conversion ± S.D. (n = 2) in % . Non-inhibited control taken as 100%. S.D. : standard deviation. NI: no inhibition

### *Inhibitory effect of the new compounds*

We investigated three picolyl (**1–3**) and three picolinylidene (**4–6**) androstene compounds (Table 1). Picolyl derivatives exerted no significant inhibition either on the  $17\alpha$ -hydroxylase or the  $C_{17,20}$ -lyase activities under our experimental conditions. Relative conversions were found between 66–100%. Elimination of the hydroxy group and introduction of a double bond in the side chain of the picolinylidene derivatives enhanced inhibitory actions. The two 3-hydroxy picolinylidene derivatives displayed moderate effect.  $IC_{50}$  values of the compounds **4** and **5** were found 38  $\mu$ M and 48  $\mu$ M against the  $17\alpha$ -hydroxylase, and 8.2  $\mu$ M and 14  $\mu$ M against the  $C_{17,20}$ -lyase, respectively. Results revealed that compound **6**, a picolinylidene derivative bearing 4-ene-3-oxo structure in its steroidal A-ring, is the best inhibitor among the tested compounds. For this compound values of  $IC_{50} = 5.9 \mu$ M for the  $17$ -hydroxylase activity and  $IC_{50} = 2.5 \mu$ M towards the  $C_{17,20}$ -lyase activity were determined.

## DISCUSSION

$P450_{17\alpha}$  lies at the crossroads of steroid hormone biosynthesis. Its inhibition reduces testosterone production and provides a feasible way for androgen withdrawal therapy. *In vitro*  $P450_{17\alpha}$  inhibition test is an essential step in the development of new antiandrogen pharmacons and it is applied furthermore in toxicology and in screening of xenobiotics or endocrine disruptors [11, 21, 29, 40].

Various laboratory techniques have been developed in the latest decades for the investigation of  $17\alpha$ -hydroxylase and  $C_{17,20}$ -lyase activities of the  $P450_{17\alpha}$ . Biotransformations are more often monitored via conversion of  $^3\text{H}$  or  $^{14}\text{C}$  labelled substrate steroids. Following the enzyme reaction labelled product steroids are isolated with chromatography and then quantified by radiometric detection [3, 6, 10–11, 21–22, 29, 40]. Other versions of the radiosubstrate method use specifically labelled substrates. Hydroxylation of the [ $17\alpha$ - $^3\text{H}$ ]Prog eliminates a tritiated water molecule [30, 39], whereas cleavage of labelled C-17 side chain of  $17\text{OH}$ -Prog produce labelled acetic acid [19, 23, 24, 28, 30]. In these techniques, enzyme activities, can be detected via measurement of labelled water or acetic acid which remains in the incubate water phase following an organic extraction. Further incubation techniques apply non-labelled substrate steroids. Enzyme products in these methods can be either isolated by high performance liquid chromatography and detected by ultraviolet spectrophotometry [20, 34–35] or can be directly quantified by specific immunoassays [9]. Technologies based on quantification of radio-labelled steroidal enzyme products, nevertheless, are acknowledged superior to other methods in sense of both precision and qualitative identification.

Here we report an improved radiosubstrate method for specific determination of the  $17$ -hydroxylase and  $C_{17,20}$ -lyase activities of  $P450_{17\alpha}$ . The newly developed technique bears numerous advantages compared to our earlier radiosubstrate methodologies [13, 37–38]. New procedures use small volume incubates, and apply simple and



efficient isolation techniques making these methods suitable for rapid and cost effective estimation of 17-hydroxylase and C<sub>17,20</sub>-lyase activities and their inhibition by test substances.

In this study, steroidal 17-picolyl and 17-picolinylidene derivatives were investigated against 17-hydroxylase and C<sub>17,20</sub>-lyase activities of P450<sub>17 $\alpha$</sub> . It has been found that the 17-picolinylidene-androstene compounds exerted stronger inhibition than their 17-picolyl counterparts. 17-Picolinylidene-androstenes bear trigonal C-17 centre, whereas this region displays tetrahedral geometry in the 17-picolyl compounds [8]. Compounds bearing heterocyclic side chain attached in trigonal geometry to the steroid C-17 are more potent inhibitors, in general, than those containing tetrahedral C-17 junction [5]. The rigid planar structure of our 17-picolinylidene groups, in particular, stabilizes the nitrogen in a position which is preferred for co-ordination to the heme of the enzyme [1]. Double bond at C17-C20 in the 17-picolinylidene compounds extends  $\pi$ -electron conjugation of the heterocyclic aromatic ring. This electron delocalization may also promote binding to the heme iron and enhance inhibitor affinity, as it is reported about several  $\Delta^{16}$  unsaturated C-17 heterocyclic steroidal compounds [5, 17, 24]. The 17-picolinylidene-androst-4-en-3-one compound (**6**) exerted effective inhibition towards both the 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase activities of the P450<sub>17 $\alpha$</sub> .

Our results demonstrate that the presented methods provide feasible ways for determination of activities and inhibitions of 17 $\alpha$ -hydroxylase and C<sub>17,20</sub>-lyase of the P450<sub>17 $\alpha$</sub> . Our findings indicate that steroidal 17-picolinylidene compounds may possibly be promising target molecules of P450<sub>17 $\alpha$</sub>  inhibitor research. Synthesis of further 17-picolinylidene steroids and investigations to explore presumed antiandrogenic properties of the new compounds are to be performed.

#### ACKNOWLEDGEMENTS

This work was supported financially by the Hungary–Serbia IPA Cross-border Co-operation Programme, project no. HURSB/1002/214/133 RECODAC. Nikolett Szabó's work was supported by a PhD Fellowship of the Talentum Fund of Richter Gedeon Plc. Authors thank the Provincial Secretariat for Science and Technological Development of the Autonomous Province of Vojvodina, grant no. 114-451-3716/2011-01. Authors are grateful to Professor Gyula Schneider and Professor János Wölfling for providing abiraterone.

#### REFERENCES

1. Ajduković, J. J., Djurendić, E. A., Petri, E. T., Klisurić, O. R., Celić, A. S., Sakač, M. N., Jakimov, D. S., Gaši, K. M. (2013) 17(E)-picolinylidene androstane derivatives as potential inhibitors of prostate cancer cell growth: antiproliferative activity and molecular docking studies. *Bioorg. Med. Chem.* 21, 7257–7266.
2. Attard, G., Reid, A. H., Yap, T. A., Raynaud, F., Dowsett, M., Settatree, S., Barrett, M., Parker, C., Martins, V., Folkard, E., Clark, J., Cooper, C. S., Kaye, S. B., Dearnaley, D., Lee, G., de Bono, J. S. (2008) Phase I clinical trial of a selective inhibitor of CYP17, abiraterone acetate, confirms that castration-resistant prostate cancer commonly remains hormone driven. *J. Clin. Oncol.* 26, 4563–4571.
3. Auchus, R. J., Lee, T. C., Miller, W. L. (1998) Cytochrome b5 augments the 17,20-lyase activity of human P450c17 without direct electron transfer. *J. Biol. Chem.* 273, 3158–3165.

4. Baston, E., Leroux, F. R. (2007) Inhibitors of steroidal cytochrome p450 enzymes as targets for drug development. *Recent Pat. Anticancer Drug Discov.* 2, 31–58.
5. Burkhart, J. P., Gates, C. A., Laughlin, M. E., Resvick, R. J. (1996) Inhibition of steroid C<sub>17(20)</sub> lyase with C-17-heteroaryl steroids. *Bioorg. Med. Chem.* 4, 1411–1420.
6. Chasalow, F. (1979) Mechanism and control of rat testicular steroid synthesis. *J. Biol. Chem.* 254, 3000–3005.
7. Djurendić, E. A., Ajduković, J. J., Sakač, M. N., Čsanádi, J. J., Kojić, V. V., Bogdanović, G. M., Penov Gaši, K. M. (2012) Synthesis and cytotoxic activity of some 17-picolyl and 17-picolinylidene androstane derivatives. *Eur. J. Med. Chem.* 54, 784–792.
8. Djurendić, E., Daljev, J., Sakač, M., Čanadi, J., Santa, S. J., Andrić, S., Klisurić, O., Kojić, V., Bogdanović, G., Djurendić-Brenesel, M., Novaković, S., Gasi, K. P. (2008) Synthesis of some epoxy and/or N-oxy 17-picolyl and 17-picolinylidene-androst-5-ene derivatives and evaluation of their biological activity. *Steroids* 73, 129–138.
9. Duc, I., Bonnet, P., Duranti, V., Cardinali, S., Rivière, A., De Giovanni, A., Shields-Botella, J., Barcelo, G., Adje, N., Carniato, D., Lafay, J., Pascal, J. C., Delansorne, R. (2003) In vitro and in vivo models for the evaluation of potent inhibitors of male rat 17 $\alpha$ -hydroxylase/C17,20-lyase. *J. Steroid Biochem. Mol. Biol.* 84, 537–542.
10. Ehmer, P. B., Jose, J., Hartmann, R. W. (2000) Development of a simple and rapid assay for the evaluation of inhibitors of human 17 $\alpha$ -hydroxylase-C(17,20)-lyase (P450c17) by coexpression of P450c17 with NADPH-cytochrome-P450-reductase in *Escherichia coli*. *J. Steroid Biochem. Mol. Biol.* 75, 57–63.
11. Fernandes, D., Bebianno, M. J., Porte, C. (2007) Mitochondrial metabolism of 17 $\alpha$  hydroxyprogesterone in male sea bass (*Dicentrarchus labrax*): a potential target for endocrine disruptors. *Aquat. Toxicol.* 85, 258–266.
12. Ferraldeschi, R., Sharifi, N., Auchus, R. J., Attard, G. (2013) Molecular pathways: Inhibiting steroid biosynthesis in prostate cancer. *Clin. Cancer Res.* 19, 3353–3359.
13. Frank, É., Mucsi, Z., Szécsi, M., Zupkó, I., Wölfling, J., Schneider, Gy. (2010) Intramolecular approach to some new D-ring-fused steroidal isoxazolidines by 1,3-dipolar cycloaddition: synthesis, theoretical and in vitro pharmacological studies. *New J. Chem.* 34, 2671–2681.
14. Gaši, K. M., Brenesel, M. Dj., Djurendić, E. A., Sakač, M. N., Čanadi, J. J., Daljev, J. J., Armbruster, T., Andrić, S., Sladić, D. M., Božić, T. T., Novaković, I. T., Juranić, Z. D. (2007) Synthesis and biological evaluation of some 17-picolyl and 17-picolinylidene androst-5-ene derivatives. *Steroids* 72, 31–40.
15. Ideyama, Y., Kudoh, M., Tanimoto, K., Susaki, Y., Nanya, T., Nakahara, T., Ishikawa, H., Yoden, T., Okada, M., Fujikura, T., Akaza, H., Shikama, H. (1998) Novel nonsteroidal inhibitor of cytochrome P450(17 $\alpha$ ) (17 $\alpha$ -hydroxylase/C17-20-lyase), YM116, decreased prostatic weights by reducing serum concentrations of testosterone and adrenal androgens in rats. *Prostate* 37, 10–18.
16. Igaz, P., Tömböl, Z., Szabó, P. M., Likó, I., Rácz, K. (2008) Steroid biosynthesis inhibitors in the therapy of hypercortisolism: theory and practice. *Curr. Med. Chem.* 15, 2734–2747.
17. Jarman, M., Barrie, S. E., Llera, J. M. (1998) The 16,17-double bond is needed for irreversible inhibition of human cytochrome P450<sub>17 $\alpha$</sub>  by abiraterone [17-(3-pyridyl)androsta-5,16-dien-3 $\beta$ -ol] and related steroid inhibitors. *J. Med. Chem.* 41, 5375–5381.
18. Jarman, M., Smith, H. J., Nicholls, P. J., Simons, C. (1998) Inhibitors of enzymes of androgen biosynthesis: cytochrome P45017 $\alpha$  and 5 $\alpha$ -steroid reductase. *Nat. Prod. Rep.* 15, 495–512.
19. Johnson, D. C. (1987) Quantitative changes in ovarian 17  $\alpha$ -hydroxylase/C17,20-lyase and aromatase activities during the estrous cycle of the hamster. *Proc. Soc. Exp. Biol. Med.* 184, 14–18.
20. Kimura, K., Itakura, Y., Goto, R., Tojima, M., Egawa, N., Yoshihama, M. (2007) Inhibition of 17 $\alpha$ -hydroxylase/C17,20-lyase (CYP17) from rat testis by green tea catechins and black tea theaflavins. *Biosci. Biotechnol. Biochem.* 71, 2325–2328.
21. Laurenzana, E. M., Balasubramanian, G., Weis, C., Blydes, B., Newbold, R. R., Delclos, K. B. (2002) Effect of nonylphenol on serum testosterone levels and testicular steroidogenic enzyme activity in neonatal, pubertal, and adult rats. *Chem. Biol. Interact.* 139, 23–41.

22. Li, J., Li, Y., Son, C., Brodie, A. M. (1995) Inhibition of androgen synthesis by 22 hydroximino-23,24-bisnor-4-cholen-3-one. *Prostate* 26, 140–150.
23. Ling, Y. Z., Li, J. S., Kato, K., Liu, Y., Wang, X., Klus, G. T., Marat, K., Nnane, I. P., Brodie, A. M. (1998) Synthesis and in vitro activity of some epimeric 20α-hydroxy, 20-oxime and aziridine pregnene derivatives as inhibitors of human 17α hydroxylase/C17,20-lyase and 5α-reductase. *Bioorg. Med. Chem.* 6, 1683–1693.
24. Ling, Y. Z., Li, J. S., Liu, Y., Kato, K., Klus, G. T., Brodie, A. M. (1997) 17-Imidazolyl, pyrazolyl and isoxazolyl androstene derivatives. Novel steroid inhibitors of human cytochrome C<sub>17,20</sub>-lyase (P450<sub>17α</sub>). *J. Med. Chem.* 40, 3297–3304.
25. Miljković, D., Gaši, K. (1982) A novel synthetic approach to 3β-acetoxy-17 picolinylidene-5-androstene-16-one. *Bull. Soc. Chim. Beograd* 47, 173–181.
26. Miller, W. L., Auchus, R. J. (2011) The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr. Rev.* 32, 81–151.
27. Moreira, V. M., Salvador, J. A., Vasaitis, T. S., Njar, V. C. (2008) CYP17 inhibitors for prostate cancer treatment—an update. *Curr. Med. Chem.* 15, 868–899.
28. Moreira, V. M., Vasaitis, T. S., Njar, V. C., Salvador, J. A. (2007) Synthesis and evaluation of novel 17-indazole androstene derivatives designed as CYP17 inhibitors. *Steroids* 72, 939–948.
29. Nishizato, Y., Imai, S., Yabuki, M., Kido, H., Komuro, S. (2010) Development of relevant assay system to identify steroidogenic enzyme inhibitors. *Toxicol In Vitro* 24, 677–685.
30. Orr, T. E., Taylor, M. F., Bhattacharyya, A. K., Collins, D. C., Mann, D. R. (1994) Acute immobilization stress disrupts testicular steroidogenesis in adult male rats by inhibiting the activities of 17 α-hydroxylase and 17,20-lyase without affecting the binding of LH/hCG receptors. *J. Androl.* 15, 302–308.
31. Owen, C. P., Shahid, I., Lee, W. Y., Ahmed, S. (2010) Synthesis and biochemical evaluation of a range of (4-substituted phenyl)sulfonate derivatives of 4-hydroxybenzyl imidazole-based compounds as potent inhibitors of 17α-hydroxylase/17,20-lyase (P45017α) derived from rat testicular microsomes. *Bioorg. Med. Chem. Lett.* 20, 5345–5348.
32. Potter, G. A., Barrie, S. E., Jarman, M., Rowlands, M. G. (1995) Novel steroidal inhibitors of human cytochrome P45017α (17α-hydroxylase-C17,20-lyase): potential agents for the treatment of prostatic cancer. *J. Med. Chem.* 38, 2463–2471.
33. Potter, G. A., Hardcastle, I. R., Jarman, M. (1997) A convenient, large-scale synthesis of abiraterone acetate [3β-acetoxy-17-(3-pyridyl)androst-5,16-diene], a potential new drug for the treatment of prostate cancer. *Organic Preparations and Procedures Int.* 29, 123–134.
34. Rotstein, D. M., Kertesz, D. J., Walker, K. A., Swinney, D. C. (1992) Stereoisomers of ketoconazole: preparation and biological activity. *J. Med. Chem.* 35, 2818–2825.
35. Sergejew, T., Hartmann, R. W. (1994) Pyridyl substituted benzocycloalkenes: new inhibitors of 17 α-hydroxylase/17,20-lyase (P450 17 α). *J. Enzyme Inhib.* 8, 113–122.
36. Sonpavde, G., Attard, G., Bellmunt, J., Mason, M. D., Malavaud, B., Tombal, B., Sternberg, C. N. (2011) The role of abiraterone acetate in the management of prostate cancer: a critical analysis of the literature. *Eur. Urol.* 60, 270–278.
37. Szécsi, M., Ondré, D., Tóth, I., Magony, S., Wölfling, J., Schneider, Gy., Julesz, J. (2010) Determination of 5α-reductase type 1 isozyme activity and its inhibition by novel steroidal oxazolines. *Acta Biol. Hung.* 61, 274–281.
38. Tóth, I., Szécsi, M., Julesz, J., Faredin, I. (1997) Activity and inhibition of 3-β-hydroxysteroid dehydrogenase/delta-5-4-isomerase in human skin. *Skin Pharmacol.* 10, 160–168.
39. Warshaw, M. L., Johnson, D. C., Azhar, S., Gibori, G. (1987) Opposite effect of human chorionic gonadotropin on placental and ovarian synthesis of androstenedione. *Endocrinology* 120, 2003–2010.
40. Ye, L., Zhao, B., Hu, G., Chu, Y., Ge, R. S. (2011) Inhibition of human and rat testicular steroidogenic enzyme activities by bisphenol A. *Toxicol Lett.* 207, 137–142.